

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

CLETHO 240 EC

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 240 g/L CLETHODIM

SOLVENT: 663g/L LIQUID HYDROCARBON

GROUP A HERBICIDE

For the control of certain grass weeds in beetroot, cabbage, canola, celery, chickpeas, cotton, faba beans, field peas, forestry, lentils, lettuce, lupins, mung beans, non-bearing fruit trees, onions, ornamentals, peanuts, potatoes and soybeans as per the Directions for Use.



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 3217 9788
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 58945/ 20L/ 0908

KENSO AGCARE CLETHO 240 EC

SELECTIVE HERBICIDE

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, or waterways with the product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers. Triple or preferably pressure rinses containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter. After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands arms and face thoroughly with soap and

water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade

Product of China



Batch No.:

Date of Manufacture:

CAUTION

KEEP OUT OF REACH OF THE CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



CLETHO 240 EC

Selective Herbicide

ACTIVE CONSTITUENT: 240g/L CLETHODIM
SOLVENT: 663g/L LIQUID HYDROCARBONS

GROUP A HERBICIDE

For the control of certain grass weeds in beetroot, cabbage, canola, celery, chickpeas, cotton, faba beans, field peas, forestry, lentils, lettuce, lupins, mung beans, non-bearing fruit trees, onions, ornamentals, peanuts, potatoes and soybeans as per the Directions for Use table

READ THIS LEAFLET BEFORE USE

APVMA Approval No: 58945/0908



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 32179788

DIRECTIONS FOR USE

RESTRAINTS: DO NOT apply without the addition of an adjuvant.

DO NOT apply to plants that are stressed by moisture or temperature extremes.

DO NOT apply if rain is expected within one hour of application.

DO NOT apply Cletho 240 EC Selective Herbicide more than once to any one crop.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING **GROUP A HERBICIDE**

Cletho 240 EC Selective Herbicide is a member of the cyclohexanedione group of herbicides. Cletho 240 EC Selective Herbicide has the inhibition of acetyl-coA carboxylase mode of action. For weed resistance management, Cletho 240 EC Selective Herbicide is a Group A Herbicide. Some naturally occurring weed biotypes resistant to Cletho 240 EC Selective Herbicide and other Group A Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Cletho 240 EC Selective Herbicide or other Group A Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Cletho 240 EC Selective Herbicide to control resistant weeds.

MIXING

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. Add spray additive and agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION

Ground Application:
Cletho 240 EC Selective Herbicide should be applied with calibrated spray equipment producing a median droplet range of 200 to 300 microns VMD. Apply in a minimum of 50 litres of water per hectare. Use 150 L/ha when spraying dense populations.

Aerial Application:

Cletho 240 EC Selective Herbicide can be applied through aircraft fitted with boom or Micron air equipment. A spray volume of 20 to 30 L/ha is recommended and equipment should be adjusted to deliver droplets in the range of 200 to 250 microns VMD. Best results will be obtained when aerial applications are made in a light crosswind. Applications should not be made during temperature inversions or in conditions of very low relative humidity. Care should be taken to avoid drift damage to adjoining grass crops.

COMPATIBILITY

Cletho 240 EC Selective Herbicide is compatible with the following grass herbicides: Quizalofop-P-ethyl, Fluazifop-P, Haloxypop-R, Sethoxydim and Propaquizafop. NOTE: In all situations where Cletho 240 EC Selective Herbicide is mixed with another grass herbicide, Cletho 240 EC Selective Herbicide should be used at the full label rate specified for the target weed species.

Cletho 240 EC Selective Herbicide may also be applied in tank mixtures with one of the following products: Esfenvalerate, Sumi-Alpha¹ Dual, Ken-Tac 100 and Omethoate.

Cletho 240 EC Selective Herbicide should not be applied in a tank mix with Legacy 500, Bentazone or Acifluorfen. Apply as separate applications. For information on compatibility of products not listed, please refer to Kenso Agcare.

CAUTION

Ornamentals and Trees - While Cletho 240 EC Selective Herbicide is generally selective to broad leaved plants (i.e. it is active against grasses), tests should always be made on a small number of plants not previously tested. Do NOT use on Gymnosperms (pines, conifers, etc) unless a prior test has been conducted to check safety on the relevant species.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do NOT apply Cletho 240 EC Selective Herbicide if wind is likely to cause drift onto susceptible crops/plants, cropping lands or pastures. Cletho 240 EC Selective Herbicide should not be applied through misting equipment or any other method likely to cause excessive drift. Care should be taken to avoid damage to adjoining native grasses or grass crops.

CROP	WEEDS CONTROLLED	RATE (mL/ha)	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Canola, Chickpeas, Faba beans, Field peas, Lentils ¹ , Lupins [#]	Annual ryegrass (<i>Lolium rigidum</i>) Annual phalaris (<i>Phalaris minor</i>)	150 to 250	2-leaf to fully tillered	Qld, NSW, ACT, Vic, Tas, SA, WA only	Always apply with D-C- Trate * at 2 L/100L or Hasten * at 1 L/100L or Kwickin* at 1L/100L or Uptake * at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides. ¹ Application up to 7-node/ early-branching crop growth stage only.
	Barley grass (<i>Hordeum leporinum</i>) Brome grass (<i>Brome diandrus</i>) Wild Oats (<i>Avena spp.</i>)	175 to 250			
	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	200* to 250 *(175 to 250 in WA only)			
	Volunteer barley (<i>Hordeum vulgare</i>)	250			
	Silver grass (<i>Vulpia bromoides</i>) – suppression only	250 to 500			
	Paradoxa grass (<i>Phalaris paradoxa</i>)	250 to 375 375			
Cotton (Qld, NSW and NT only), Peanuts, Mung beans, Soybeans	Barnyard grass (<i>Echinochloa spp.</i>), Blown grass (<i>Agrostis aveacea</i>), Crowsfoot grass (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass (<i>Sorghum halepense</i>), Summer grass (<i>Digitaria spp.</i>), Volunteer sorghum (<i>Sorghum spp.</i>)	250 to 375 375	2- to 5- leaf stage 5 leaf to fully tillered	Qld, NSW, ACT, Vic, WA, NT only	Always apply with D-C- Trate at 2L/100L or Hasten at 1L/100L or Kwickin at 1L/100L or Uptake at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides.
	Paradoxa grass (<i>Phalaris paradoxa</i>)	175 to 250	2- to 5- leaf stage	Qld, NSW, ACT only	
	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	200* to 250* (175 to 250 in WA only)	5 leaf to fully tillered	Qld, NSW, ACT, Vic, WA, NT only	
	Volunteer barley (<i>Hordeum vulgare</i>)	250			
	Silver grass (<i>Vulpia bromoides</i>) – suppression only (not Qld, WA)	250 to 500	5 leaf to fully tillered	Qld, NSW, ACT, Vic, WA, NT only	
	Winter grass (<i>Poa annua</i>)	500			

CROP	WEEDS CONTROLLED	RATE (mL/ha)	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Beetroot, Cabbage, Celery, Lettuce, Potatoes, Onions	Barnyard grass (<i>Echinochloa spp.</i>) Blown grass (<i>Agrostis aveacea</i>), Cowsfoot grass (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Paradoxa grass (<i>Phalaris paradoxa</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass (<i>Sorghum halepense</i>), Summer grass (<i>Digitaria spp.</i>), Volunteer sorghum (<i>Sorghum spp.</i>)	250 to 375 375	2- to 5- leaf stage 5-leaf to fully tillered	All States	Always apply with D-C-Trate at 2 L/100L or Hasten at 1L/100L or Kwickin at 1L/100L or Uptake at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. Use a spray volume of 150 L/ha when spraying dense grass populations.
	Annual ryegrass (<i>Lolium rigidum</i>), Annual phalaris (<i>Phalaris minor</i>)	150 to 250	2-leaf to fully tillered	All States	Always apply with D-C-Trate at 2L/100L or Hasten at 1L/100L or Kwickin at 1L/100L or Uptake at 500mL/100L spray volume.
	Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus diandrus</i>), Wild Oats (<i>Avena spp.</i>)	175 to 250	5 leaf to fully tillered	Qld, NSW, ACT only	
	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	200* to 250* (175 to 250 in WA only)			
	Volunteer barley (<i>Hordeum vulgare</i>)	250	5 leaf to fully tillered	Qld, NSW, ACT, Vic, WA, NT only	
	Silver grass (<i>Vulpia bromoides</i>) – suppression only (not Qld, WA)	250 to 500			
	Winter grass (<i>Poa annua</i>)	500	5 leaf to fully tillered	Qld, NSW, ACT, Vic, WA, NT only	

CROP	WEEDS CONTROLLED	RATE (mL/ha)	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Forestry, non-bearing fruit trees and ornamentals	Annual ryegrass (<i>Lolium rigidum</i>), Annual phalaris (<i>Phalaris minor</i>), Barley grass (<i>Hordeum leporinum</i>), Barnyard grass (<i>Echinochloa spp.</i>), Blown grass (<i>Agrostis aveaeae</i>), Brome grass (<i>Bromo diandrus</i>), Crowsfoot (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Paradoxa grass (<i>Phalaris paradoxa</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass (<i>Sorghum halepense</i>), Silver grass (<i>Vulpia bromoides</i>)-suppression only (not Qld, WA), Summer grass (<i>Digitaria spp.</i>), Volunteer sorghum (<i>Sorghum spp.</i>), Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer barley (<i>Hordeum vulgare</i>), Winter grass (<i>Poa annua</i>)	500	2-leaf to fully tillered	All States	Apply after plants have recovered from transplant shock and are showing signs of active growth. Always with D-C-Trate at 2L/100L or Hasten at 1L/100L or Kwickin at 1L/100L or Uptake at 500 mL/100L spray volume. Use a spray volume of 150 L/ha when spraying dense grass populations. Do NOT use on Gymnosperms (pines, conifers, etc) unless a prior test has been conducted to check safety on the relevant species. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for particular species.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST:

CANOLA, COTTON, PEANUTS, CHICKPEAS, FABA BEANS, FIELD PEAS, FORESTRY, LENTILS, LUPINS, MUNG BEANS, SOYBEANS: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

BEETROOT, CABBAGE : DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

ONIONS : DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.

LETTUCE, POTATOES : DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST.

CELERY : DO NOT APPLY LATER THAN 9 WEEKS BEFORE HARVEST.

GRAZING:

CANOLA, PEANUTS, CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS, MUNG BEANS, SOYBEANS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION.

COTTON : DO NOT GRAZE OR CUT COTTON FORAGE OR STUBBLE FOR STOCK FEED.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well-ventilated area out of direct sunlight. Protect from frost. Triple, or preferably, pressure rinse containers before disposal. Add rinsing to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non-return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter.

After each use of the product, please ensure that the non-return valve, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If products in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 32179788

APVMA Approval No: 58945/0908