

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KENSO AGCARE

KEN-ESTER LV 680

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 680 g/L 2, 4-D
present as the 2-ETHYLHEXYL ESTER

GROUP I HERBICIDE

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the Directions for Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 3217 9788
www.kenso.com.au

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 Litres APVMA Approval No.: 65224/ 50453

KENSO AGCARE KEN-ESTER LV 680 SELECTIVE HERBICIDE

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KEN-ESTER LV
680

Selective Herbicide
ACTIVE CONSTITUENT: 680 g/L 2,4-D present as the 2-ETHYLHEXYL ESTER

GROUP **1** HERBICIDE

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the Directions for Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

IMPORTANT: READ THE LEAFLET BEFORE USE

APVMA Approval No: 65224/50453



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 3217 9788

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycle or designated collection point.

If not recycling, break, crush or puncture container and deliver empty packaging for appropriate disposal to an approved waste management facility. Empty containers and product should not be burnt.

110L Mini Bulk Returnable Container

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the non return valve or the security seal. DO NOT contaminate the drum with water or any foreign matter.

After each use of the product, please ensure that the non return coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.

Refillable containers (1000L only):

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty containers fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Kenso Corporation (M) Sdn. Bhd.
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 3217 9788

APVMA Approval No: 65224/50453

DIRECTIONS FOR USE:

RESTRAINTS: AVOID spraying if rain is likely within 6 hours or if strong winds prevail. DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat, Barley	Refer to Weed Table	Vic only	210-800mL	CROP STAGES: ALL CEREALS Apply at tillered to boot stages. (WA, Vic only). Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. (NSW, ACT only). Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling of the head at the top of the main stem. (Old only). Apply from completion of tillering to early jointing stage. (SA, Tas only).
		SA only	230-800mL	
		QLD, NSW, ACT only	410-800mL	
		Tas only	620-800mL	
		WA only	800mL	
Triticale		SA, Qld, NSW, ACT only	410-800mL	
		Vic only	210-800mL	
Cereal Rye		NSW, Qld, ACT only	410-800mL	
		Vic only	210-800mL	
Sugar cane		Qld only	1.1-2.4L	Post-emergence
Stubble/ Fallow Spray Prior to Direct Drilling or Sowing. Winter Cereals, Grain legumes (Peanuts Old only), Canola		All States	210-800mL/ha	Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as Roundup [®] Max, Credit [®] plus Bonus [®] , Ken-up 450 CT, Glyphosate CT, Paraken 250 or Speedy 250. Select appropriate rate from the weed table. For skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
Harvest Aid or Salvage Spray	Broadleaf Weeds	All States	1.7L	Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For desiccation of green matter, estimate harvest date and apply spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results. NB. Where thistles are tall and branching above the crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages. Spraying may increase seed contamination of harvest by accelerating maturity. Do not use with undersown legumes that have not set seed.
-Winter Cereals	Refer to Weed Table			
Potatoes Pre-harvest Preparation	Broadleaf Weeds such as Clover, Variegated Thistle & Cruciferous weeds	Vic, Tas only	1.1-2.4L	Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30cm in height. For boom spraying apply at least 100 litres of spray mixture per hectare. If grasses such as Rye grass and winter grass are also present add Amitrole T Herbicide.

2. PASTURES, NON-AGRICULTURAL, INDUSTRIAL

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Improved Pasture containing Clovers	Refer to Weed Table	NSW, Tas, SA, Qld, ACT only	410-800mL	Clover must be well covered by the grass or extensive damage may result.
		NSW, Tas, SA, WA, Qld, ACT only	800mL-4.7L	
		Vic only	800mL -6.6L 70-620mL	
Pastures - non legumes, Rights of way & Industrial	Charlock, Clover, Medics, Mustards, Paterson's Curse, Saffron, Slender, variegated and Spear Thistles, Turnip weed, wild Radish, wild Turnip	NSW, ACT only	800mL-1.5L (Aerial application)	Apply to young, actively growing weeds. Sowing: Do not sow pasture seed for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
			800mL - 1.1L (Ground Application)	
			3.3-4.7L (Aerial or Ground)	
			As above plus Propron or Roundup [®] Max or Credit [®] plus Bonus [®] or Glyphosate CT or Weedmaster [®] Duo	
			All of Above plus grasses	
Pastures - Direct Drilling or Surface Sowing	As Above plus: Capeweed, Wireweed, Storksbill/Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles		800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	
			800mL	

3. SPOT SPRAYING

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Spot Spraying (All Situations)	Refer to Weed Table	All States	1/100 th of rate on Weed Table per 10L water per 100m ²	Apply through Knapsack. Thorough wetting of weed is essential.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION. IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15TH APRIL TO 15TH SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD (WHP): PASTURES, CEREALS CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

APPLICATION INFORMATION

This product may be used in either high and low volume sprays. Just pour into water and stir. BOOM SPRAYING - Use 30-100 litres water per hectare. AERIAL SPRAYING - Use 10-90 litres water per hectare. NOTE: Refer to the Department of Agriculture/ Primary Industries in your state for the current restricted spraying areas.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. Do not use wooden spray vats as they can not be cleaned. Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

COMPATIBILITY

This product can be tank mixed with 500g/L Dicamba, Ken-Chlor 750, Paraken 250, Propron[®], Atrazine 900 WG Herbicide, Credit[®]/Bonus[®] Roundup[®] Max, Roundup[®] CT, Ken-up 450 CT, Ken-Met 600, Ken-Gran 750, Speedy 250[®], Weedmaster Duo.

WEED TABLE

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds may only be suppressed. E.g. Rumex spp. (docks) and Polygonum spp. (wiredraw climbing buckwheat) are killed to ground level only.

APPLICATION RATE PER HECTARE

Weeds Controlled	CROP							PASTURES - NON LEGUMES		Critical Comments
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, Tas, SA, Qld, WA, ACT only		
Amaranthus spp	-	-	-	800mL	-	-	-	-	-	Spray when buds forming or early flowering
Angled Onion	-	-	-	-	-	-	3.3L	0.8-1.7L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Apple of Sodom	-	-	-	-	-	-	-	2.9-3.3L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Bathurst Burr	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	-	Spray before flowering. DO NOT cultivate these infestations
Black Knapweed	-	-	-	-	-	-	3.3L	-	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Buffalo Burr	-	-	-	-	-	-	-	800mL - 1.1L (not Qld & WA)	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Callitrop	-	-	-	620-800mL	-	-	1.7-3.3L	-	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Cape Tulip	-	-	-	-	-	1.1L	3.3L	1.7-3.3L	-	Spray before flowering.
Capeweed	800m L	800m L	800m L	530-800mL	-	-	-	2.5-3.3L	-	Spray up to rosette stage
Charlock	410m L	410m L	800m L	410mL	-	-	-	800mL	-	Spray up to rosette stage.
Clover	-	-	-	620-800mL	-	-	-	-	-	
Colocynth	-	-	-	-	-	-	3.3L	-	-	Spray at seedling stage only
Deadnettle	-	-	-	800mL	-	-	-	-	-	
Devil's Claw	-	-	-	800mL	-	-	1.3L	-	-	Spray prior to pods forming.
Dock	800m L	800m L	-	-	800m L	800mL	-	1.7-2.5L	-	Spray at rosette stage to kill top growth only
Fat Hen	-	-	-	410-800mL	-	-	-	-	-	
Flatweed	-	-	-	800mL	-	-	-	2.5-3.3L	-	
Fumitory (red)	-	800m L	-	800mL	-	-	-	2.5-3.3L	-	Spray up to rosette stage
Fumitory (white)	800m L	410m L	-	800mL	-	-	-	2.5-3.3L	-	Spray up to rosette stage
Galvanised Burr	-	-	-	-	-	-	4.7L	4.7L (not Qld & WA)	-	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	800mL	-	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.3-5.2L	-	-	Spray before flowering
Hogweed, Wireweed	800m L	800m L	-	800mL	800m L	-	-	1.1-1.7L (not SA)	-	Spray up to rosette stage
Hoary Cress.	-	800m L	800m L	800mL	-	-	1.7-3.3L	1.7-2.1L	-	Spray from late rosette to pre-flowering.
Horehound (seedlings)	-	800m L	-	-	-	840mL	-	1.7-3.3L	-	Late Autumn to early Spring
Ironweed, Com Gromwell	-	-	-	800mL	-	-	-	1.1-1.7L	-	
Khaki Weed	-	-	-	-	-	-	-	800mL-1.1L (Not SA)	-	Spray young seedlings only
Lincoln Weed	-	800m L	-	-	-	-	-	-	-	Autumn spray before sowing improves control.

Weeds Controlled	CROP						PASTURES –NON LEGUMES		
	Vic	SA	Vic	SA	Vic	SA	Vic	SA	
London Rocket	-	-	-	-	-	570mL	-	1.6-2.5L (WA only)	
Lupins	800mL	-	-	410mL-800mL	-	-	-	-	Spray up to rosette stage.
Melilotus/ Hexham Scant	800mL	800mL	-	-	800mL	-	-	1.1-1.7L	Spray up to rosette stage.
Melons – camel, paddy	-	-	-	410mL-800mL	-	-	-	-	
Mustards	330mL	230-800mL	800mL	410mL-800mL	620mL	620mL	3.3L	1.7-2.5L	Spray up to rosette stage.
Mexican Poppy	-	-	-	800mL	-	840mL	-	800mL-1.1L (1.1-1.5L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	800mL	620mL	-	-	800mL-1.1L	Spray active seedlings only
Muskweed	800mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	800mL	-	-	-	-	
Noogoora Burr	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	Spray seedling to pre-flowering
Nut Grass	-	-	-	-	-	-	3.3-5.2L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	-	800mL	-	840mL	1.7-3.3L	800mL-1.7L(1.1-1.5L WA only)	Spray seedling to rosette stage
Poppy Wild	410mL	-	-	-	-	-	-	2.1-2.9L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.3L	3.3L	Spray at rosette to cabbage stage.
Rapeseed	800mL	-	-	410mL-800mL	-	-	-	-	Spray up to rosette stage.
Rapistrum	-	-	-	-	-	570mL	-	840mL (WA only)	
Rough Poppy	-	410mL	-	410mL-800mL	-	-	-	800mL	Spray young seedlings only.
St. John's Wort	-	-	-	-	-	-	3.3-5.2L	3.3-4.7L	Spray before flowering. Spray before plants are 40cm high.
Safflower	-	-	-	410mL-800mL	-	-	-	-	
Sand Mustard/ Sand Rocket	-	-	-	-	-	-	3.3L	-	Spray before flowering
Shepherd's Purse	-	-	-	800mL	-	-	-	-	
Silver Leaf Nightshade	-	-	-	-	-	-	3.3L	-	Spray at flowering. Followland: controls top growth only
Skeleton Weed	-	800mL	-	800mL	-	-	3.3L	1.1-1.7L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	800mL	800mL	-	-	-	-	2.1-2.5L	
Stinging Nettle	800mL	-	-	-	-	-	-	-	Spray up to rosette stage
Stinkwort	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	Spray younger plants. Use higher rate as plants mature.
Storkbill/ Erodium	-	-	-	800mL	-	-	-	-	
Sunflower Seedlings	800mL	-	-	410mL-800mL	620mL	-	-	-	Spray multiple leaves.
Thistles:	-	-	-	-	-	-	-	-	
-Golden	-	-	-	-	-	-	3.3L	3.3L	Spray at rosette stage
-Nodding	-	-	-	-	-	-	3.3L	1.1-1.7L	Spray rosette stage to pre-flowering
-Saffron	620mL	800mL	-	410mL-800mL	800mL	800mL	800mL-1.7L	800mL-2.5L	Spray up to rosette stage.
-Sheep	-	-	-	-	-	840mL	-	840mL-3.3L (WA only)	
-Slender, Shore	-	-	-	800mL	800mL	-	1.7-3.3L	800mL-3.3L	Spray at rosette stage.
-Soldier	-	-	-	-	-	-	3.3L	-	Spray at rosette stage
-Spear	800mL	-	-	800mL	-	-	800mL-2.5L	1.1-2.1L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures)
-Stemless	-	-	-	-	-	-	3.3L	2.5-3.3L	Spray rosette stage to flowering.
-St. Barnaby's	-	-	-	-	-	-	-	1.1-1.7L	
-Star	-	-	-	800mL	-	-	1.7-3.3L	-	Spray seedling to rosette stage. Use higher rate as plants mature.
-Variegated	-	-	-	800mL	410mL-800mL	620mL	-	800mL-3.3L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	-	-	410mL-800mL	-	-	3.3L	800mL-1.7L	Spray at seedling stage
Tree Hogweed	800mL	-	-	-	-	-	-	-	Spray up to rosette stage
Turnip Weed	-	410mL	-	410mL-800mL	410mL	620mL	-	800mL	Spray seedlings only
Vetches/Tares	800mL	620mL	800mL	-	-	-	-	-	
Wards Weed	-	410mL	-	-	-	-	-	-	Spray at seedling stage
Wild cabbage	800mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Wild Garlic	-	-	-	-	-	-	6.6L	-	Suppresses aerial growth only
Wild Mignonette	-	-	-	-	-	840mL	3.3L	-	Spray at rosette stage
Wild Mustard	-	-	-	-	-	570mL	-	1.6-2.3L (WA only)	
Wild Radish	800mL	800mL	800mL	410mL-800mL	800mL	570mL	-	800mL (840mL WA only)	Spray up to rosette stage
Wild Sage	-	-	-	-	-	-	-	2.5-3.3L	
Wild Teasel	-	-	-	-	-	-	1.7-3.3L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	210mL	230mL	800mL	410mL-800mL	-	400mL	-	800mL (840mL WA only)	Spray up to rosette stage.

Plant back days for Kenso Agcare Ken-Ester LV 680 Selective Herbicide

Crop	Rates		
	Up to 510mL/ha	510mL-1L/ha	1-1.6L/ha
Balansa Clover	7	7	10
Barley%	1	1	3
Chickpeas#	7	14	21
Cotton	10	14	21
Faba beans	7	7	10
Field peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medics	7	7	10
Marbon beans	7	7	10
Navybean	10	10	14
Oats	3	3	7
Perennial ryegrass	7	7	10
Persian clover	7	7	10
Phalaris	7	7	10
Canola/rapeseed #	14	21	28
Rice	7	7	14
Safflower#	7	14	21
sorghum@	3	7	10
Soybean	14	14	21
Sub-clover	7	7	10
Sunflower@	7	10	14
Triticale%	1	3	7
Vetch	7	7	10
Wheat%	1	3	7
White Clover	7	7	10

IMPORTANT:WHEN APPLIED TO DRY SOILS AT LEAST 15MM (½inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and Triticale.

In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15mm.

@ In Central Queensland, when using 830mL /ha or less of Ken-Ester LV 680, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.

+ In WA the Plant Back Period for Lupins at all rates is 28 days.

RESISTANT WEEDS WARNING

GROUP	HERBICIDE
I	

Kenso Agcare Ken-Ester LV 680 Selective Herbicide is a member of the Phenoxy group of herbicides Ken-Ester LV 680 has the Disruptors of plant cell growth mode of action. For weed resistance Ken-Ester LV 680 is a group I herbicide. Some naturally-occurring weed biotypes resistant to Ken-Ester LV 680 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Ken-Ester LV 680 or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use. Kenso Corporation accepts no liability for any losses that may result from the failure of Ken-Ester LV 680 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Legume Tolerance: If clovers are present, care should be taken to ensure that they are reached the 3-4 leaf stage before spraying. Rates above 410mL of this product per hectare will destroy most clovers, whilst Lucerne and medics are susceptible at any strength.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. In particular, avoid spray drift and vapour movement onto susceptible crops such as: cotton, tobacco, tomatoes, flowers, vegetables, vines, fruit trees, legume crops and pastures, oilseed crops or other susceptible plants and trees (eg. Kurrajongs, Eucalypts and Belahs).

DO NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site. DO NOT apply with smaller than coarse to very coarse spray droplets according to the BCPC/ASAE S572 definition for standard nozzles.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any times as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.