


POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Pollux



ACTIVE CONSTITUENT : 250 g/L FLUTRIAFOL

GROUP **C** FUNGICIDE



For the control of certain fungal disease on wheat, barley and canola when mixed with fertilizer or applied as a foliar spray as specified in the Direction for Use table.

**IMPORTANT: READ THE ATTACHED
LEAFLET BEFORE USE**

CONTENTS:

20 Litres

APVMA Approval No.:

60027/20/0606



Pollux

20L

KENSO AGCARE POLLUX FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinses the containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. If product in eyes wash it out immediately with water. If product on skin immediately wash area with soap and water. Wash hands after use.

When opening the container and preparing spray wear:

- cotton overalls, buttoned to the neck and wrist;
- a washable hat;
- elbow length PVC, nitrile or neoprene gloves;
- goggles; and
- disposable fume mask.

When using the prepared spray wear:

- cotton overalls, buttoned to the neck and wrist;
- a washable hat;
- elbow length PVC, nitrile or neoprene gloves; and
- goggles.

After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (TEL: 131126). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data which is available from the supplier.

CONDITIONS OF SALE

"Kenso Corporation (M) Sdn. Bhd." ('Kenso') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add to or alter these conditions.

In a Transport Emergency
Dial **000**
Police or Fire Brigade

Batch No:

Date of Manufacture:

Product of China



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Pollux

Fungicide

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP **C** FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertilizer or applied as a foliar spray as specified in the "Direction for Use" table.

DIRECTIONS FOR USE LEAFLET

APVMA Approval No: 60027/0606



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone 07 3217 9788

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"Kenso Corporation (M) Sdn. Bhd." ("Kenso") shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add to or alter these conditions.

In a Transport Emergency
Dial **000**
Police or Fire Brigade

Batch No:

Date of Manufacture:



Kenso Corporation (M) Sdn. Bhd.
3C / 59 Oxford Street
Bulimba QLD 4171
Phone: 07 3217 9788

APVMA Approval No: 60027/0606

DIRECTIONS FOR USE:

A. Pollux Fungicide as a fertilizer treatment

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments																												
Canola	Blackleg (<i>Leptosphaeria macularis</i>)	400mL	4 weeks grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as guide.																												
				Fertilizer Rate Per hectare	Volume of Product per 100kg of fertilizer (mL)																											
				60	667*																											
				70	571																											
				80 500 100 400 120 333																												
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	200 or 400mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertilizer in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as guide.																												
	Septoria Tritic blotch (<i>S. tritici</i>), take-all (<i>Geaumannomyces graminis</i>)	400mL																														
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	200 or 400mL		<table border="1"> <thead> <tr> <th>Fertilizer Rate Per hectare</th> <th>Volume of Product per 100kg of fertilizer (mL)</th> <th>200mL Rate</th> <th>400mL Rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>400</td> <td>800*</td> <td></td> </tr> <tr> <td>60</td> <td>333</td> <td>667*</td> <td></td> </tr> <tr> <td>70</td> <td>286</td> <td>571</td> <td></td> </tr> <tr> <td>80</td> <td>250#</td> <td>500</td> <td></td> </tr> <tr> <td>100</td> <td>200#</td> <td>400</td> <td></td> </tr> <tr> <td>120</td> <td>167#</td> <td>333</td> <td></td> </tr> </tbody> </table> <p>*Application volumes over 600mL per 100kg of fertilizer may adversely effect the flow rate and handling characteristic of dusty/poorly granulated fertilizers. #Where application volume is low, product may be diluted with water to improve coverage.</p>	Fertilizer Rate Per hectare	Volume of Product per 100kg of fertilizer (mL)	200mL Rate	400mL Rate	50	400	800*		60	333	667*		70	286	571		80	250#	500		100	200#	400		120	167#	333	
Fertilizer Rate Per hectare	Volume of Product per 100kg of fertilizer (mL)	200mL Rate	400mL Rate																													
50	400	800*																														
60	333	667*																														
70	286	571																														
80	250#	500																														
100	200#	400																														
120	167#	333																														

B. Pollux Fungicide Applied as a Foliar Spray

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	250-500mL Ground Applications: Apply in 100-150mL water plus 200mL "Ken-Wett 1000" per 100L of spray solution.	7 weeks harvest or grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf rust (<i>Puccinia striiformis</i>) Septoria nodorum blotch (<i>Septoria nodorum</i>) Septoria Tritic blotch (<i>S. Tritici</i>)	Aerial Application: Apply in a minimum of 15L water plus 200mL "Ken-Wett 1000" per 100L of spray solution.		Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe diseases persist during flowering, use a single application at the higher rate or two applications 3-4 weeks apart at the low rate.
Barley	Powdery mildew (<i>Erysiphe graminis</i>)		10 weeks harvest or grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

A. Pollux Fungicide Applied in-furrow as a fertilizer

TREATMENT:

CANOLA, WHEAT and Barley: DO NOT GRAZE OR CUT FOR STOCK FOR 4 WEEKS AFTER SOWING.

B. Pollux Fungicide Applied as a Foliar Spray

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION.

BARLEY – 10 WEEKS

WHEAT – 7 WEEKS

DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION.

BARLEY – 10 WEEKS

WHEAT – 7 WEEKS

GENERAL INSTRUCTIONS

Fertilizer Application:

Ensure that treated fertilizer is dry before sowing with the crop.

Sowing: Treated fertilizer should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertilizer may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the corrected rates of fertilizer and fungicide are delivered. DO NOT mix seed with treated fertilizer.

Foliar Application:

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

Fungicide Resistance Warning

GROUP	C	FUNGICIDE
-------	---	-----------

Pollux Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Pollux Fungicide is a Group C Fungicide. Some naturally occurring individual fungi resistant to Pollux Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Pollux Fungicide and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Corporation (M) Sdn. Bhd. accepts no liability for any losses that may result from the failure of Kenso Agcare Pollux Fungicide to control resistant fungi. Resistant should not be assumed without first reviewing the method of application, the coverage and the time of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers.

Triple or preferably pressure rinses containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any foreign matter.

After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.