

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
AG CARE

**Thidiazuron**

**500 SC**

**Cotton Defoliant**

ACTIVE CONSTITUENT : 500 g/L THIDIAZURON

For the defoliation of cotton prior to harvest


**IMPORTANT: READ THE ATTACHED  
LEAFLET BEFORE USE**



**CONTENTS:**

**5 Litres**

**APVMA Approval No:  
56469/5/1003**



Thidiazuron 500SC 5L

Kenso Corporation (M) Sdn Bhd 3C / 59 Oxford Street Bulimba QLD 4171 Phone (07) 3217 9788

## KENSO AGCARE THIDIAZURON 500 SC COTTON DEFOLIANT

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT allow spray drift to contact trees or crops other than cotton, as this product may injure or defoliate other crops, particularly citrus, grapevines and lettuce.

To help reduce drift damage potential:

- DO NOT apply Kenso Agcare Thidiazuron 500SC by air when citrus in flush is within 8 km downwind from the point of application.
- DO NOT apply Kenso Agcare Thidiazuron 500SC by air when within 800 m of lettuce.

In addition:

- DO NOT plant lettuce or carrots in cotton fields where Kenso Agcare Thidiazuron 500SC has been applied within the previous 9 months.
- DO NOT plant legumes, lucerne, other root crops or leafy vegetables (besides those noted above) in cotton fields where Kenso Agcare Thidiazuron 500SC has been applied within the previous two months.
- DO NOT plant small grains, sorghum or corn in cotton fields where Kenso Agcare Thidiazuron 500SC has been applied within the previous two weeks.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

*For refillable containers (110L) :* Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves. After each day's use, wash gloves and contaminated clothing. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

### EXCLUSION OF LIABILITY

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

In a Transport  
Emergency Dial  
000  
Police or Fire Brigade

Batch No.:

Date of Manufacture :



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**KENSO**  
agcare

**Thidiazuron**

**500 SC**

**Cotton Defoliant**

**ACTIVE CONSTITUENT: 500 g/L THIDIAZURON**

For the defoliation of cotton prior to harvest

**READ THIS LEAFLET BEFORE USING THIS PRODUCT**

**APVMA Approval No: 56469/1003**



**Kenso Corporation (M) Sdn Bhd**  
3C / 59 Oxford Street  
Bulimba QLD 4171  
Phone (07) 3217 9788

**DIRECTIONS FOR USE** (For use in NT, Qld, NSW, WA only)

**Restraints**

DO NOT apply Kenso Agcare Thidiazuron 500SC when night temperatures fall below 12° C.

DO NOT apply Kenso Agcare Thidiazuron 500SC following application of sodium chlorate, magnesium chlorate or diquat.

**GENERAL INSTRUCTIONS**

Kenso Agcare Thidiazuron 500SC is an odourless, non-corrosive defoliant/harvest aid. After treatment with Kenso Agcare Thidiazuron 500SC, leaves fall while still green so that there is minimal dry leaf trash to contaminate fibre or interfere with harvesting. Maximum defoliation occurs 7 to 21 days following application of Kenso Agcare Thidiazuron 500SC. Good control of regrowth is seen when Kenso Agcare Thidiazuron 500SC is used at rates over 100 mL/ha. Defoliation will be slower under average conditions, or when the crop is not physiologically mature.

**Note**

Rainfall within 24 hours after application will reduce the effectiveness of Kenso Agcare Thidiazuron 500SC; Retreatment may be necessary. Defoliation of crops with moderate to high levels of nitrogen remaining in the soil are more prone to leaf freezing than crops that have depleted soil nitrogen.

Tank mixes of Kenso Agcare Thidiazuron 500SC with organophosphates may form a residue in application equipment. While still fresh and moist, this residue is easily removed by flushing the entire system with a commercial tank cleaner.

Do NOT allow the spray solution to dry in the application equipment. Immediately following application, flush and rinse equipment (mix tanks, pumps, transfer lines, application tanks, sumps, booms, nozzles and all related equipment) thoroughly with detergent and water.

Should any residue of Kenso Agcare Thidiazuron 500SC remain in the sprayer system, it may be released during subsequent applications of other products causing damage to crops.

**Export of Treated Produce**

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Kenso Agcare Thidiazuron 500SC. If you are growing produce for export, please check with Kenso Corporation (M) Sdn Bhd for the latest information on MRLs and import tolerances BEFORE using Kenso Agcare Thidiazuron 500SC.

CROP/SITUATION	DEFOLIATION CONDITIONS	RATE/HA	CRITICAL COMMENTS
Cotton Defoliation - first or second spray	<b>IDEAL</b> Conditions are hot to warm (30 to 35°C daytime temperature) and the crop is ready to be defoliated with natural senescence already occurring. Nitrogen and water are almost depleted and there is little or no top growth. A good fruit load is set.	50 to 100 mL plus 2.0 L Ampol D-C-TRON® Cotton Spray Oil	Apply the first spray to cotton, which has been grown under good (non-stressed) conditions and is physiologically mature i.e., when the top harvestable boll is mature. Application made prior to boll maturity may result in yield and quality reductions. <ul style="list-style-type: none"> <li>• Use the lower rates of 50 to 75 mL/ha only under warm early season conditions.</li> <li>• Use the higher rate for denser canopies or to provide a faster response.</li> <li>• A second spray 7 to 10 days later of either Kenso Agcare Thidiazuron 500SC or an alternative harvest aid may be required.</li> </ul>
	<b>GOOD</b> Conditions are warm (25 to 30°C daytime temperature) and the crop is ready to be defoliated. Nitrogen and water are low and there is little or no new growth and a good fruit load is set.	100 to 150 mL plus 2.0 L Ampol D-C-TRON Cotton Spray Oil	Apply as above, but use these rates under good conditions.
	<b>AVERAGE</b> Conditions are: (i) warm but night temperatures are dropping to 12°C (not below) and the crop is physiologically ready to defoliate or; (ii) temperature is still hot/warm but the crop is not physiologically ready to senesce.	150 to 200 mL plus 2.0 L Ampol D-C-TRON Cotton Spray Oil	Apply as above, but use these rates under average conditions.
	<b>COLD</b> Night temperatures fall below 12°C and/or mean day/night temperatures fall below 18°C.	-	DO NOT apply Kenso Agcare Thidiazuron 500SC under cold conditions.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD**

Cotton harvesting - NOT REQUIRED WHEN USED AS DIRECTED.

Grazing/Cotton Trash - DO NOT GRAZE TREATED AREAS OR FEED COTTON TRASH TO LIVESTOCK.

## Mixing

Add the required amount of Kenso Agcare Thidiazuron 500SC straight into the partly filled spray tank or mixing vat with agitator running, followed by Ampol D-C-TRON Cotton Spray Oil, then add the rest of the water. Keep mixture agitated while spraying.

Use only freshly prepared sprays. **Do NOT store spray mixture overnight.**

When tank mixing with emulsifiable concentrates (ECs), add the Kenso Agcare Thidiazuron 500SC to the ¼ filled spray tank or mixing vat and agitate thoroughly. With agitator running, then add the EC product, followed by the Ampol D-C-TRON Cotton Spray Oil. Keep mixture agitated while spraying.

## Application

Apply as a uniform spray ensuring thorough coverage of the whole plant.

## Equipment

### Aircraft

Apply by aircraft in 20 to 40 L/ha spray mixture and ensure thorough coverage.

A steady cross wind is recommended for best penetration. Do not exaggerate swath width.

### Ground Sprayers

Standard boom sprays are recommended and must be fitted with mechanical or by-pass agitation. Use sufficient water to obtain complete coverage of all leaves, ie at least 90 to 100 litres of water per hectare. Good coverage should be obtained with three or more nozzles per row depending upon crop height and canopy density. A nozzle configuration including droppers will give maximum coverage in dryland crops with particular care being taken to ensure thorough but even coverage. Overdosing on some plant parts, eg., nozzle overlap along the sides of plants may cause localised leaf freezing. If band spraying dryland crops, care should be taken when calculating the actual applied rate of Kenso Agcare Thidiazuron 500SC per sprayed hectare.

## Compatibility

Kenso Agcare Thidiazuron 500SC is compatible in tank mixtures with ethephon, dimethoate and parathion.

CAUTION: When tank mixed with Roundup® CT (glyphosate), a higher proportion of cotton leaf may be retained (freezing). Do NOT use this mixture on stressed cotton, particularly drought affected cotton.

## PRECAUTIONS

Do not use human flagging unless flaggers are protected by engineering controls such as vehicles with cabs.

### Re-entry Period

Do not allow entry into treated areas until spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT allow spray drift to contact trees or crops other than cotton, as this product may injure or defoliate other crops, particularly citrus, grapevines and lettuce.

To help reduce drift damage potential:

- D
- NOT apply Kenso Agcare Thidiazuron 500SC by air when citrus in flush is within 8 km downwind from the point of application.

- D
- NOT apply Kenso Agcare Thidiazuron 500SC by air when within 800 m of lettuce.

In addition:

- D
- NOT plant lettuce or carrots in cotton fields where Kenso Agcare Thidiazuron 500SC has been applied within the previous 9 months.

- D
- NOT plant legumes, lucerne, other root crops or leafy vegetables (besides those noted above) in cotton fields where Kenso Agcare Thidiazuron 500SC has been applied within the previous two months.

- D
- NOT plant small grains, sorghum or corn in cotton fields where Kenso Agcare Thidiazuron 500SC has been applied within the previous two weeks.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For refillable containers (110L) :** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves. After each day's use, wash gloves and contaminated clothing. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26).

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

## EXCLUSION OF LIABILITY

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

D-C-TRON® is a Registered Trademark of Ampol Australia Pty. Ltd.

Roundup® is a Registered Trademark of Monsanto Company USA.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade
---

NRA Approval No: 56469/0903



Kenso Corporation (M) Sdn. Bhd.  
3C / 59 Oxford Street  
Bulimba QLD 4171