

Cletho 240

Selective post-emergent grass herbicide

About Cletho 240

The active constituent of Cletho 240 'Clethodim' is renowned throughout Australia for its excellent activity on numerous grass weeds including one of Australia's most devastating weeds, Annual Ryegrass.

The "DIM" chemistry of Cletho 240 provides a valuable tool in the control of herbicide resistant Ryegrass.

Effective weed control of grass weeds in broadleaf crops is an important tool in Australian Broadacre rotations.



Product Details:

Product:

Cletho 240

Active Constituent:

240g/L Clethodim

Herbicide Group:

GROUP A HERBICIDE

Pack Sizes:

4 x 5L Carton, 20L, 110L

Primary Uses:

For the control of certain grass weeds in canola, field peas faba beans, lentils, soybeans, chickpeas, mungbeans, peanuts and various horticultural and forestry uses.

APVMA Approval Number:

58945/5/0307

Best Use Tips:

- Do not apply without addition of an adjuvant
- Do not apply to plants stressed by moisture or temperature extremes
- Do not apply Cletho 240 more than once to any crop
- Do not apply if rain is expected within one hour of application
- Adhere to the withholding periods stated on the label
- Do not graze or cut for stock feed for 8 weeks after application
- Apply when target weeds are healthy and growing with vigour
- Spray weeds when they are young and small.





Grass weeds controlled by Cletho 240:

Winter weeds:

- Barley Grass, Brome Grass, Annual Phalaris, Annual Ryegrass, Paradoxa Grass, Volunteer Barley, Volunteer Oats, Volunteer Wheat, Wild Oats, Winter Grass, Suppression of Silver Grass.

Summer weeds:

- Barnyard Grass, Blown Grass, Crowsfoot Grass, Feathertop Rhodes Grass, Liverseed Grass, Red Sprangletop Grass, Seedling Johnston Grass, Summer Grass, Volunteer Sorghum.

Guaranteed to perform:

Our products are guaranteed to perform when used as directed or your product is replaced.

Kenso Agcare produces and distributes a wide range of quality crop protection products for use in Australian farming systems.

As a division of Kenso Corporation, we have over 30 years experience in agricultural chemical manufacture and formulation. From our three modern production sites we are able to efficiently cater to the seasonal demand peaks of the Australian market.

We also have a dedicated product development and quality control chemistry team which provides the capability to design robust product formulations for Australian conditions.

In line with our motto “**Better Crop Production for a Growing Population**” Kenso has the vision of being recognised as a world class supplier of crop protection products for Australian growers.

Major Broadacre Uses (see the full label for horticultural and forestry uses)

CROP	WEEDS CONTROLLED	RATE (mL/ha)	STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
Canola, Chickpeas, Faba Beans, Field peas, Lentils ¹ , Lupins [#]	Annual ryegrass (<i>Lolium rigidum</i>) Annual phalaris (<i>Phalaris minor</i>)	150 to 250	2-leaf to fully tillered	QLD, NSW, ACT, VIC, TAS, SA, WA only	Always apply with D-C-Trate* at 2 L/100L or Hasten* at 1 L/100L or Kwickin* at 1 L/100L or Uptake* at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides.
	Barley grass (<i>Hordeum leporinum</i>) Brome grass (<i>Brome diandrus</i>) Wild oats 9Avena spp.)	175 to 250			
	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	200* to 250* (175 to 250 in WA only)			
	Volunteer barley (<i>Hordeum vulgare</i>)	250	NSW, ACT, VIC, TAS only		
	Silver grass (<i>Vulpia bromoides</i>) – suppression only	250 to 500			
	Paradoxa grass (<i>Phalaris paradoxa</i>)	250 to 375 375	2-to-5-leaf stage 5 leaf to fully tillered	QLD, NSW, ACT only	
Cotton (QLD, NSW and NT only), Peanuts, Mung beans, Soybeans.	Barnyard grass (<i>Echinochloa</i> spp), Blown grass (<i>Agrostis aveacea</i>), Crowsfoot grass (<i>Eleusine indica</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass (<i>Sorghum halepense</i>), Summer grass (<i>Digitaria</i> spp), Volunteer sorghum (<i>Sorghum</i> spp).	250 to 375 375	2-to-5-leaf stage 5 leaf to fully tillered	QLD, NSW, ACT, VIC, WA, NT only	Always apply with D-C-Trate* at 2 L/100L or Hasten* at 1L/100L or Kwickin* at 1L/100L or Update* at 500mL/100L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides.

*Please note: This is a Cletho 240 label extract only. The full version of the label can be found at www.kenso.com.au