

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## Vega 250 EC

### Fungicide

ACTIVE CONSTITUENT : 250 g/L DIFENOCONAZOLE  
SOLVENTS : 500 g/L HYDROCARBON LIQUID  
200 g/L N-METHYL PYRROLIDONE

GROUP **C** FUNGICIDE



For the control of target spot of potatoes and  
tomatoes, leaf blight of carrots, leaf spot  
diseases of bananas and husk spot  
on Macadamias as per the  
Directions for Use table

**IMPORTANT: READ THE ATTACHED LEAFLET  
BEFORE USE**

### CONTENTS:

20 Litres

APVMA Approval No.:  
61153/20L/0807



9 328666 001409

Vega 250

20L

## KENSO AGCARE VEGA 250 EC FUNGICIDE

### STORAGE AND DISPOSAL

Store in closed container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty container and product should not be burnt.

### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin, avoid contact with eyes and skin. When opening the container and preparing spray wear:

- cotton overalls buttoned to the neck and wrist,
- a washable hat,
- elbow-length butyl rubber gloves and
- face shield or goggles.

When using the prepared spray wear:

- cotton overalls buttoned to the neck and wrist,
- a washable hat and
- elbow-length butyl rubber gloves.

If product gets in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed **DO NOT** induce vomiting. Give a glass of water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

### CONDITIONS OF SALE

"Kenso Corporation (M) Sdn. Bhd." ('Kenso') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add to or alter these conditions.

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade

Batch No :

Date of Manufacture:



## CAUTION

KEEP OUT OF REACH OF THE CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
CORPORATION

## Vega 250 EC

### Fungicide

ACTIVE CONSTITUENT: 250 g/L DIFENOCONAZOLE  
SOLVENTS: 500 g/L HYDROCARBON LIQUID  
200 g/L N-METHYL PYRROLIDONE

GROUP **C** FUNGICIDE

For the control of target spot of potatoes and tomatoes, leaf blight of carrots, leaf spot diseases of bananas and husk spot on Macadamias as per the Directions for Use table

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No: 61153/0807



Kenso Corporation (M) Sdn Bhd  
3C / 59 Oxford Street  
Bulimba QLD 4171  
Phone (07) 3217 9788

## DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT apply more than 6 applications of this product per season. The effect of Vega 250 could be diminished if rain falls within 2 hours of application.

Crop	Disease Controlled	State	Rate/ha	WHP	Critical Comment
<b>Bananas</b> Including bananas interplanted with avocados	Yellow Sigatoka ( <i>Mycosphaerella musicola</i> ), Black Sigatoka ( <i>Mycosphaerella fijiensis</i> )	Qld, NSW, NT only	<b>Ground Application</b> 400mL + 3 to 5L of water miscible oil, in a convenient volume of water.  <b>Aerial Application</b> 100mL + 3 to 5L of a water miscible oil, in a minimum of 30L of water	1 day	<b>This use is subject to a DMI anti-resistant strategy.</b> <b>Ground Application:</b> Apply by misting machine or airblast sprayer. Use rates towards the higher end of the range where weather conditions favour diseases or where equipment or terrain does not permit thorough spray coverage of all foliage. <b>NSW, Sth Qld: Ground and aerial application:</b> Commence spraying at the start of the summer rainy season. Apply a maximum of 5 Group C sprays per season. Apply at least 2 consecutive sprays of Vega 250 at 21 to 28 day intervals before applying further treatments of a protectant fungicide from a different activity group. <b>NT, Nth Qld: Ground and aerial applications:</b> Commence spraying at the start of the wet season. Apply a regular schedule of protectant sprays. When the disease potential is high, apply at least 2 but no more than 3 consecutive sprays of Vega 250 at 14 to 21 day intervals before applying further treatments of a protectant fungicide from a different activity group. Apply a maximum of 6 Group C sprays per season. Continue with treatments using an alternative protectant fungicide for the remainder of the season. Use the lower rate of oil in Nth Qld. DO NOT apply during July, August, September and October.
<b>Carrots</b>	Leaf Blight ( <i>Alternaria dauci</i> and <i>Cercospora carotae</i> )	All states	300mL or 500mL	7 days	<b>This use is subject to an AVCARE anti-resistance strategy.</b> Use in a protectant program commencing before disease occurs. Limit the use of Group C fungicides to periods when conditions favour disease development. Apply the 300mL rate at 7 day intervals and the 500mL rate at 10 to 14 day intervals. Use the 500mL rate when conditions favour high disease pressure. Apply no more than 2 consecutive sprays of Vega 250 alone. Then apply a recommended non Group C fungicide or tank mix Vega 250 with a recommended protectant fungicide for at least 1 application before continuing with Vega 250 alone. Good crop hygiene will aid disease control.

### 110L Mini Bulk Returnable Container (110L only)

Store the original sealed drum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any foreign matter.

After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed **DO NOT** induce vomiting. Give a glass of water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### CONDITIONS OF SALE

"Kenso Corporation (M) Sdn. Bhd." ('Kenso') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add or alter these conditions.

In a Transport Emergency  
Dial **000**  
Police or Fire Brigade



Kenso Corporation (M) Sdn. Bhd.  
3C / 59 Oxford Street  
Bulimba QLD 4171  
Phone 07 3217 9788

not a trademark of Kenso Corporation (M) Sdn Bhd

## DIRECTIONS FOR USE – continued

Crop	Disease controlled	State	Rate/ha	WHP	Critical Comments
<b>Macadamia nuts</b>	Husk Spot ( <i>Pseudocercospora macadamiae</i> )	QLD, NSW, NT only	<b>Dilute spraying</b> 50mL/100L water  <b>Concentrate spraying</b> Refer to the Application section	-	Apply by dilute or concentrate spraying equipment. <b>Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.</b> Use in a protectant fungicide program containing fungicides from different chemical groups. Commence fungicide applications at nut set (approximately late September) and continue the fungicide program until late December, with applications at 3 to 4 week intervals. Apply a maximum of 2 applications of Vega 250 per season. Ensure thoroughly spray coverage. Add Agravit® spray adjuvant at a rate of 10mL/100L.
<b>Potatoes</b>	Target Spot/ Early Blight ( <i>Alternaria solani</i> )	ALL States	300mL or 500mL	7 days	<b>This use is subject to an AVCARE anti-resistance strategy.</b> Use in a protectant program commencing when rows meet and before disease occurs. Limit the use of Group C fungicides to periods when conditions favour disease development. Apply the 300mL rate at 7 day intervals and the 500mL rate at 10 to 14 day intervals. Use the 500mL rate when conditions favour high disease pressure. Apply no more than 2 consecutive sprays of Vega 250 alone. Then apply a recommended non Group C fungicide or tank mix Vega 250 with a recommended protectant fungicide for at least 1 application before continuing with Vega 250 alone. Good crop hygiene will aid disease control. Vega 250 does not control late blight, Phytophthora infestans. If this disease is a problem, mix Vega 250 with an appropriate fungicide for control of this disease and observe the spray interval for that product.
<b>Tomatoes</b>	Target Spot ( <i>Alternaria solani</i> )	All States	300mL or 500mL	3 days	<b>This use is subject to an AVCARE anti-resistance strategy.</b> Use in a protectant program commencing soon after transplanting. Limit the use of Group C fungicides to periods when conditions favour disease development. Apply the 300mL rate at 7 day intervals and the 500mL rate at 10 day intervals. Apply no more than 2 consecutive sprays of Vega 250 alone. Then apply a recommended non Group C fungicide or tank mix Vega 250 with a recommended protectant fungicide for at least 1 application before continuing with Vega 250 alone. Good crop hygiene will aid disease control.

NOT TO BE USED FOR ANY PRUPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:	
Bananas:	DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.
Potatoes, Carrots:	DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.
Tomatoes:	DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.
Avocados, Macadamias:	NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Fungicide Resistance Warning

GROUP C FUNGICIDE

Vega 250 EC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Vega 250 EC Fungicide is a Group C fungicide. Some naturally occurring individual fungi resistant to Vega 250 EC Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Vega 250 EC Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Kenso Corporation (M) Sdn Bhd accepts no liability for any losses that may result from the failure of Vega 250 EC Fungicide to control resistant fungi.

Mixing

Add the required amount of Vega 250 directly to the spray tank and mix well.

Application

Thorough coverage is essential for effective results.  
Aerial application: Use 20 to 30L water / ha.  
Ground application: Use at least 100L water / ha.

Macadamia nuts: Apply by high volume (dilute) sprayer or by concentrate sprayer.

Dilute spraying: Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

Concentrate spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: for example 2000L/ha
2. Your chosen concentrate spray volume: for example 500L/ha
3. The concentration factor in this example is: 4 x (ie 2000L ÷ 500L = 4)
4. If the dilute label rate is 50mL/100L, then the concentrate rate becomes 4 x 50, that is 200mL/100L of concentrate spray.

The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Compatibilities

This product is compatible with methamidophos, dimethoate, methidathion, Kenso Agcare Kensban 500, pirimicarb, permethrin, Kenso Agcare Ken-Tac 100, Kenso Agcare Kencozeb 750 formulations, chlorothalonil, procymidone, iprodione, mancozeb, Kendon High K, urea, EC formulations of synthetic pyrethroids and organophosphate insecticides and formulations of copper oxychloride.

PRECAUTION:

RE-ENTRY PERIOD

**DO NOT** enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist and butyl rubber gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied to plants any time.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to a aquatic organisms. **DO NOT** spray within 5 m downwind of water bodies. **DO NOT** apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses. **DO NOT** contaminate dams, waterways or drains with the product or its container.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.